


Spatial Smoothing and Small-Sample Bias in Mortality Forecasting



Jack C. Yue

National Chengchi University

Date: Sept. 9, 2026

Email: csyue@nccu.edu.tw




Global Life Prolonging

- ❑ Life expectancy continues to rise, occasionally affected by factors such as disease & war.
 - COVID-19 reduced life expectancy by 1.5 years.
- ❑ Taiwan's Life expectancy continues to increase.
 - Annual increment appears to remain unchanged.
 - Mortality models are often used to predict life expectancy.

Note: Lee-Carter model is a fine choice.



Taiwan's Annual Increment Remains Unchanged





Lee-Carter Model

□ Lee and Carter (1992) proposed

$$\ln(m_{xt}) = \alpha_x + \beta_x \kappa_t + \varepsilon_{xt}$$

where $\alpha_x \rightarrow$ Age-specific average mortality

$\beta_x \rightarrow$ Age-specific mortality improvement

$\kappa_t \rightarrow$ Time-varying mortality index


Note: SVD, Approximation, & Poisson (or MLE)
Estimation.

Sample Size and Estimation


- ❑ Variance of estimates is inversely proportional to the sample size.
 - Parameter estimation may also be biased for the small populations.
- ❑ Applying Lee-Carter model to small populations:
 - α_x estimates are under-biased,
 - β_x estimates can be over- or under-biased in younger ages.





Variance decreases with sample size



α_x Estimates for the Lee-Carter Model



β_x Estimates of the Lee-Carter Model





Bias Reduction in α_x & β_x

- ❑ The low number of deaths leads to high sampling variability in mortality rates.
- α_x is the average of $\log(\text{mortality})$ & the bias is most significant at about ages 0 & 10.
- The bias of β_x is most obvious around ages 0, 10, & 20.

Note: The most effective way to reduce the bias of parameter estimates is to increase sample size.



Modification of Lee-Carter Model

- ❑ Modifications of Lee-Carter model are proposed to deal with small populations
 - Poisson Lee-Carter model (Brouhns et al., 2002)
 - Multiple populations (Li & Lee, 2005)
 - Bayesian Lee-Carter (Czado et al., 2005)
 - Graduation+Lee-Carter (Wang et al., 2018)
 - Spatial Lee-Carter (Leong & Yue, 2026)


Note: These methods can be seen as increasing the sample size.

Study Objective


- ❑ Mortality data in age and time format can be viewed as two-dimensional coordinates.
 - Enlarge the sample size by including nearby cells.
- ❑ Spatial modification of Data Augmentation.
 - Partial SMR smooths mortality rates using the reference population in 1-dim.
 - Treat age $\times$ time as 2-dim coordinates and apply GWR to increase the effective sample size.

Residuals of LC model (Japan)

Japan Male (LC Residuals)



Japan Female (LC Residuals)



Partial Standard Mortality Ratio

□ Lee (2003) used SMR to smooth mortality rates based on a large reference area.

$$v_x = u_x^* \times \exp \left(\frac{d_x \times \hat{h}^2 \times \log(d_x / e_x) + (1 - d_x / \sum d_x) \times \log(\text{SMR})}{d_x \times \hat{h}^2 + (1 - d_x / \sum d_x)} \right)$$
$$\text{SMR} = \frac{\sum_x d_x}{\sum_x n_x \cdot u_x^*}, \quad \hat{h}^2 = \max \left(\frac{\sum \left((d_x - e_x \times \text{SMR})^2 - \sum d_x \right)}{\text{SMR}^2 \times \sum e_x^2}, 0 \right)$$

→ n_x & d_x : Numbers of people and deaths aged x

→ u_x^* : Mortality rates aged x in the large population ◦

GWR (Geographically Weighted Regression)


- (GWR) is a modelling technique designed to deal with spatial non-stationarity, i.e., the mean values vary by locations.

$$Y_i = \beta_{i0} + \sum_{k=1}^p \beta_{ik} x_{ik} + \varepsilon_i$$

Note: Determine how to implement GWR and select the optimal bandwidth.

Bandwidth Choice Alters GWR Estimates



$$y_i = \sum_j \boldsymbol{\beta}_{ij} (u_i, v_i) X_{ij} + \boldsymbol{\varepsilon}_i$$




Computer Simulation Setting

- ❑ Mortality rates follow Lee–Carter model and parameters are estimated from Taiwan male mortality data, 1992–2020.
- Simulate no. of deaths using Poisson distribution.
- Partial SMR reference: Pool data across all years within each age group (1992–2020).
- 5-Age groups: 0–4, 5–9, 10–14, ..., 80–84 (17).
- Population sizes: 10,000, 20,000, 50,000, 100,000, 200,000, and 500,000.

α_x Estimates of Partial SMR



β_x Estimates of Partial SMR



Partial SMR & Parameter Estimation



□ Partial SMR shows improvement on estimating mortality rates when the population size is 20,000~500,000.

→ Significant improvement on the bias of α_x estimate.



→ Little improvement on the bias of β_x estimate.

Note: Variances of parameter estimates are also smaller.

α_x Estimates of GWR



β_x Estimates of GWR



GWR & Parameter Estimation


- ❑ GWR has little improvement on mortality & parameter estimation, although the variances of α_x & β_x estimates are smaller.
- ❑ Why does GWR perform poorly?
 - Does not increase the actual number of deaths
 - May oversmooth real mortality patterns
 - Sensitive to bandwidth selection
 - Reduces variance more than bias

Proposed Spatial modification


- Can GWR provide additional improvement after Partial SMR smoothing?
 - Use the partial SMR to reduce small-sample bias and then apply GWR to further reduces the remaining local variability
- The bias of α_x & β_x estimates of GWR+Partial SMR do not shown improvement.

α_x Estimates of GWR & GWR+Partial SMR

GWR




PSMR + GWR




β_x Estimates of GWR & GWR+Partial SMR

GWR



PSMR + GWR



Adjustments for Proposed Approach

□ Problem identified

- Partial SMR effectively reduces the bias of α_x .
- GWR may oversmooth local mortality patterns, with limited bias reduction.
- Performance of GWR is sensitive to bandwidth.

□ Proposed adjustment:

- The key is to reduce bias first, then control local variability.

Conclusion

- Small populations induce substantial estimation bias
 - The bias is particularly serious for α_x and the bias of β_x is more evident at ages 0, 10, & 20.
- GWR alone is not sufficient to solve the small-population bias problem.
 - Partial SMR is effective for bias reduction and GWR mainly reduces variance, not bias.



Future Study

- Improved spatial modification
 - Combine Partial SMR with other spatial model
- Optimal bandwidth selection
 - Elliptical weighted window
 - Age-time dependent weighting/flexible bandwidth
- Empirical application
 - Apply the proposed method to Taiwan's small-area mortality data

Take-home Message

◆ Increasing the effective sample size can reduce small-population bias.

◆ Study notes:

Partial SMR → Effective for reducing bias

GWR → Effective mainly for reducing variance

Note: Future direction → Develop a spatial method that can improve both bias and variance.



References

- Brouhns, Denuit, and Vermunt (2002), A Poisson Log-bilinear Approach to the Construction of Projected Lifetables, *Insurance: Mathematics and Economics*.
- Czado, Delwarde, and Denuit (2005), Bayesian Poisson log-bilinear mortality projections, *Insurance: Mathematics and Economics*.
- Leong and Yue (2026), A Spatial Approach to Model Mortality Rates, *Journal of Forecasting*.
- Li and Lee (2005), Coherent Mortality Forecasts for a Group of Populations: An Extension of the Lee–Carter Method, *Demography*.
- Wang, Yue, and Chong (2018), Mortality Models and Longevity Risk for Small Populations, *Insurance: Mathematics and Economics*.
- Wiśniowski, Smith, Bijak, Raymer, and Forster (2015), Bayesian Population Forecasting: Extending the Lee-Carter Method, *Demography*.
- Yue, Wang, and Wang (2021), Using Graduation to Modify the Estimation of Lee-Carter Model for Small Populations, *North American Actuarial Journal*.

Thank you
for your Attention!

