

Modelling Healthy Life Expectancy at Neighbourhood Level

Andrew J.G. Cairns

Director, Duncan Laboratory for Insurance Data Science
Heriot-Watt University, Edinburgh

Longevity 21, Rome, September 2026



- Motivation and objectives
- Data: life, health, socio-economic
- Modelling of neighbourhood mortality and health rates
- Results
- Discussion: problems with HLE

Motivation and Objectives

- **Inequality:** *systematic, avoidable and unfair differences in mortality and health outcomes and metrics*

Main reason:

prevalence of *causal risk factors* varies between groups (e.g. smoking)
+ access to healthcare and related services

- **Short-term aim:**

To develop *customised* mortality and health indices for Scotland

Longer term: UK-wide indices (especially mortality)

- How much **inequality** is there in mortality and health rates at different ages?
- Which **socio-economic factors** are most strongly associated with mortality and health outcomes?

Mortality-Related Data: Neighbourhoods

- Small neighbourhoods:
 “Data Zones”: DZ (equivalent to LSOAs in England and Wales)
- $L = 6976$ DZs
- Average size ≈ 800 persons
- DZs $i = 1, \dots, L$,
 single years ($t = 2011-2019$),
 single or 5-year age groups, x :
 - Deaths: $D(i, t, x)$ (5-yr age groups) (by normal place of residence)
 - Exposures: $E(i, t, x)$ (single ages)
- Many potential predictive variables, $\Rightarrow$ vector $Z(i)$

- National death rates (all t and x)

$$m(t, x) = \frac{D(t, x)}{E(t, x)} \quad (\text{single ages})$$

- Neighbourhood death rates ($i = 1, \dots, L$)
 - neighbourhood death rates: $m(\mathbf{i}, t, x)$
 - Question: how to model $m(i, t, x)$ relative to national $m(t, x)$?
- Predictive variables $Z(i)$ plus $m(t, x) \longrightarrow m(i, t, x)$

Methodology: General Approach

- Over a limited age range (e.g. 70-89); and
- Over a limited range of years (e.g. 2011-2019):


neighbourhood i ,
$$m(i, t, x) = m(t, x)F(i)$$

- $F(i)$ = neighbourhood-specific *relative risk* due to socio-economic and other non-spatial characteristics, $Z(i)$
- Estimate $F(i)$ using **Local Linear Regression** (variant of LOESS)

Choice of Predictive Variables: what do we include in $Z(i)$


- **Scottish Index of Multiple Deprivation (SIMD):**
 - Employment deprivation
 - Education deprivation: Proportion of the population with no qualifications
 - Crime
- **Census data:**
 - Communal establishments: proportion of the population aged 65+ in a communal establishment (mostly care homes in the 65+ group)
 - Proportion married: ages 35+
- **2011-2019 population data:**
 - Old-age dependency ratio: $\# 65+ / \# 20-64$; males and females separately

Socio-Economic Relative Risk, $F(i)$: Significant Inequality



- High levels of inequality
[Higher levels of inequality than England.]
- Inequality diminishes with age
- Age 40-59 main predictors:
unemployment, married, education
- Age 70-89 main predictors:
care homes, education, dependency ratio
- Females slightly less spread out
High correlation between male and female
relative risks

Geographic Spread of Inequality: Males 60-79; Central Scotland




Decile 1: high mortality,

Decile 10: low mortality

Life Expectancies

- End users understand life expectancies better than relative risks
- Here: Period life expectancies
(i.e. based on mortality rates for a given group of years: 2017-2019)
- Challenges and assumptions:
 - LE calculations assume that there is no migration between areas as people get older
 - Relative risks (RR) by neighbourhood i
Modelling $\Rightarrow$ age groups 30-49; ...; 70-89
Need $RR(i, x)$ by single ages 0 to 110
Interpolate RR between 40, 50, 60, 70 and 80
RR tapers to 1 at age 100
Extrapolate back to age 0
- Combine $m(i, x) = RR(i, x)m_{NAT}(x) \Rightarrow LE(0)$ and $LE(x)$ for higher ages

Life Expectancies at Age 0, $LE(0)$: Males versus Females



- Axes: range = 22 on both axes
- Males: 64.1 to 83.1, StDev= 3.7
- Females: 69.8 to 86.0, StDev= 2.5
- Takeaway:

Most deprived areas:
female-male gap ≈ 8 years


Least deprived areas:
female-male gap ≈ 2 years

Healthy Life Expectancy (HLE)

- Good health $\leftrightarrow$ poor health $\leftrightarrow$ dead
- LE measures average years spent in both good and poor health
- **How many years are spent in good health?**
- HLE calculation relies on **self-assessment** of “general health”
- Census 2022 and Health Survey:
How is your health in general?
Possible answers: Very good; good; fair; bad; very bad
- $\pi(x)$ = proportion at exact age x who report that they are in **good or very good health**
- $m(y)$ = death rate at age y
 $S(x) = \exp[-\sum_{y=0}^{x-1} m(y)]$ = *period* survival function age 0 to age x

Sullivan Method for HLE

$$HLE(0) \equiv \frac{1}{2}\pi(0)S(0) + \sum_{x=1}^{\infty} \pi(x)S(x)$$



- *Period* HLE
- $HLE(0)$ = national HLE
- $HLE(i,0)$ = data zone i HLE using $\pi(i, x)$ and $S(i, x)$
- As with $LE(i,0)$, $HLE(i,0)$ assumes no migration between subgroups
- Different from LE: HLE is based on subjective responses

Data for estimating HLE at the data zone level

- Five health states:
very good; good; fair; bad; very bad
- Health Survey:
too small to estimate health rates accurately at the data zone level
- Census 2022:
Pro: reasonably full coverage, so data-zone analysis is theoretically possible
Con: 2022 $\Rightarrow$??? more people reporting poor health post-Covid
Con: no annual updating
Con: males and females combined (due to be separated)

Data for estimating HLE at the data zone level (cont.)

- Census outputs: headcounts by data zone subdivided into

Males and females combined

5 health states $\times$ 6 age groups (A)

0-15; 16-24; 25-34; 35-49; 50-64; 65+

- $GH(i, A)$ = number in Good or VG health in age group A in data zone i
- $P(i, A)$ = total persons in all states of health
- $\pi(i, A) = GH(i, A)/P(i, A)$ empirical proportion in good health, age group A

Estimating HLE at the data zone level

- Model:
Use the empirical $\pi(j, A)$ to estimate

$$\hat{\pi}(i, A) \equiv f(A, Z_H(i))$$

where $Z_H(i)$ = vector of socioeconomic predictive variables

- Use local linear regression (LLR) again to estimate the $\hat{\pi}(i, A)$
- Age group $\hat{\pi}(i, A) \rightarrow$ piecewise logit-linear $\hat{\pi}(i, x)$ for single ages x
- Combine with data-zone period survival curve $\hat{S}(i, x)$

$$HLE(i, 0) \equiv \frac{1}{2}\hat{\pi}(i, 0) + \sum_{x=1}^{\infty} \hat{\pi}(i, x)\hat{S}(i, x)$$


Predictive Variables for Health versus Mortality

Current variable selections:

Mortality	Health
Communal Establishment 65+ Employment Deprivation Education: Low (SIMD) Married Old-age Dependency Ratio Crime	Communal Establishment 65+ Employment Deprivation Education: Low (Census) Married Old-age Dependency Ratio Education: High (Census) UK born


Life Expectancy versus Healthy Life Expectancy from Age 0

Life Expectancy v Healthy Life Expectancy
By Data Zone for Males




- High correlation between LE and HLE
(Strong correlation between $S(i, x)$ and $\pi(i, x)$)
- Highest LE $\Rightarrow$ 8 years in "poorer" health
(females 9)
- Lowest LE $\Rightarrow$ 23 years in "poorer" health
- One year reduction in LE $\Rightarrow$ 1.75 reduction in HLE
(females 2.35)

The Index of Scottish Life Expectancy Males Healthy Life Expectancy From Age 0



(Healthy) Life Expectancy: Females vs Males



- **Life Expectancy:**

Gradient > 1

$\Rightarrow$

Males much more variable than females

- **Healthy Life Expectancy:**

Gradient ≈ 1

BUT: conclusions limited by the data:

Persons only; not M and F

Alternatives to Healthy Life Expectancy


$$\text{Healthy Life Expectancy: } HLE(i, 0) = \sum_{x=0}^{\infty} \hat{\pi}(i, x) \hat{S}(i, x) - \frac{1}{2} \hat{\pi}(i, 0)$$

$$\text{Age Standardised Health Rate: } ASHR(i) = \frac{\sum_{x=0}^{\infty} \hat{\pi}(i, x) S(x)}{\sum_{x=0}^{\infty} S(x)}$$

where $S(x)$ = national survival curve (or other standard population)

- HLE has an implicit assumption:
individuals stay in the same data zone for life
- ASHR does not require this assumption
it just gives a measure of the average health across all ages
- Formulas are very similar:
 - HLE uses custom weights
 - ASHR uses standardised weights

Comparison of HLE by Data Zone With Age Standardised Health Rate



- Very high correlation between HLE and ASHR
- Less than 100% correlation: differences in the profile of $S(i, x)$ versus national $S(x)$ period survival curves

Problems with HLE

- Health is on a sliding scale
not 0/1 like mortality
- Health is also multidimensional
- Health status is **subjective** ⇒
 - Perception of what is *good/very good* health varies with age
Mental benchmark for *good health* falls with age
 - Possible bias by socio-economic or cultural group e.g.
more deprived ⇒ ? relatively pessimistic ⇒ “poorer” health
less deprived ⇒ ? relatively optimistic ⇒ “better” health
 - Census: head of household might be responding on behalf of others
- Interpreting trends is hard!
E.g. mental health more likely now to be taken into account when self-assessing health status

Problems with HLE (cont.)

- Non-response bias in the health survey and Census 2022
Less likely to respond: lower SE group; lower education; poor health
and, non-response bias is getting worse
- Our data: Census 2022 $\Rightarrow$
potential Covid impact on health rates (real and subjective)

Possible improvements:

- ONS are looking at the use of administrative health data to replace survey data
 - More objective and better coverage
 - But relies on equal access to and willingness to access healthcare
 - How do you calculate an **objective score** for each individual?
 - Score $<$ **threshold** $\Rightarrow$ good or very good health
Calibrate this threshold to approximate old HLE's

Thank You!

E: A.J.G.Cairns@hw.ac.uk

W: www.macs.hw.ac.uk/~andrewc/ARCresources

THE Index of Scottish Life Expectancy – THISLE

